

09 Motion in Fields review questions


Part A: Projectile motion:

1. A stone is thrown from a hand at a velocity of 24ms^{-1} and an elevation of 40° to the horizontal. The stone is 2m above the ground when released. Ignore air resistance.
 - a. Calculate the horizontal and vertical components of velocity.
 - b. Calculate the time taken for the stone to reach the top of its flight.
 - c. Calculate the total flight time for the stone.
 - d. Calculate the horizontal distance travelled by the stone.
 - e. Calculate the velocity of the stone just before landing.
 - f. Sketch the flight path of the stone with arrows drawn to scale to represent the velocity of the stone at the start, high point and landing.
2. A bullet is fired horizontally from a gun at 500ms^{-1} directly at a cross on a target. If the target is 200m away how far below the cross will the bullet hit?

Part B: Gravitational field, potential and energy and orbital motion

Mass of Earth = $6 \times 10^{24}\text{kg}$, Mass of Moon = $7.3 \times 10^{22}\text{kg}$


1. A satellite is in orbit 42,000 km above the centre of the Earth.
 - a. Calculate the gravitational potential at this orbit height explaining why it is negative.
 - b. If the satellite has a mass of 150kg calculate its gravitational potential energy.
 - c. Calculate the gravitational field strength at this point and hence state the centripetal acceleration the satellite undergoes.
 - d. Hence calculate the orbit period of this satellite in days.
2. Derivation of the formula for escape velocity for a planet of mass M, radius r:
 - a. State the energy transfer involved as an object moves completely away from a planet due only to its velocity.
 - b. State the total energy and object has when infinitely far from any other mass and when travelling at an infinitesimal velocity.
 - c. State the total energy the object must have just after launch.
 - d. Hence derive the formula for escape velocity.
3. The distance from the Earth to the moon is $3.8 \times 10^8\text{m}$. A point r at a distance of $3.8 \times 10^7\text{m}$ from the moon in a direction directly towards the Earth is shown on the potential energy diagram opposite:




- a. From the graph alone state the approximate value of the gravitational field strength at this point.
- b. Calculate the gravitational potential due to the moon and the Earth combined at this point.
- c. Calculate the flight velocity required for an object, whose rockets will stop working at a distance of 7000km from the centre of the Earth to reach, this point in space.

Part C: Electric field, potential and energy

1. How much work is done to bring a charge of 1C to within 1×10^{-6} m of a charge of 10C?
2. How much work is done to bring a charge of 1C to within 1×10^{-6} m of a charge of -10C?
3. The graph below shows the variation of electric potential with distance near a point charge.
 - a. Use the graph to estimate the field strength at a distance of 1m.
 - b. Hence calculate the size of the charge.



4. Sketch the equipotential surfaces and electric field lines for:
 - a. A point positive charge
 - b. Two point negative charges separated by a small gap.



5. The image shows electric field lines that are made visible by observing tracks of charged particles moving through a liquid. **4cm separates the plates.**
 - a. If the potential difference between the plates is 6V calculate the force experienced by an ion of charge e.
 - b. How much work is done moving the charge all the way from one plate to another?
 - i. in electron volts.
 - ii. in Joules

<http://www.physics.upenn.edu/undergraduate/undergraduate-physics-labs/experiments/electric-field-and-electric-potential>